



Curzate® M-8

Fungicide

KEEP OUT OF REACH OF CHILDREN, DOMESTIC ANIMALS, FOOD AND MEDICATIONS.

USE PRECAUTIONS AND WARNINGS:

Protective Equipment: Read the instructions carefully. Evacuate the area to be people who do not have the proper protective equipment. Do not expose workers or other persons to come into direct contact or indirect (drift) with the spray. Avoid creating dust.

Applicators must wear: long sleeved shirt, long trousers (overalls), shoes and socks, gloves and chemical-type goggles / mask. People who mix and load the product, and clean the units must be used in addition to an apron of chemical.

For aerial applications, the flaggers must be protected inside booths or fold and shelter from the spray.

Do not eat, drink or smoke during the preparation and product application.

Wash clothing and protective equipment with hot soapy water separately from other laundry.

Discard clothing and other absorbent materials that have been seriously contaminated with this product. Do not reuse.

All users should: Wash hands before eating, smoking, drinking, chewing gum, tobacco or using the toilet.

Remove clothing immediately if the product passes or protective equipment after using the product, wash well and wear clean clothes. Before removing gloves wash thoroughly with soap and water.

DIRECTIONS FOR USE:

CROP	DISEASE	RATE	INTERVAL OF APPLICATION
Potato <i>Solanum tuberosum</i>	late blight	2-3 kg / ha	3 to 5 days
Tobacco <i>Nicotiana tabacum</i>	Blue Mold	2-3 kg / ha	3-5 days depending the incidence of fungus and moisture
TOMATO <i>Lycopersicon esculentum</i>	Late blight	2-3 kg / ha	3 to 5 days
CUCURBITS MELON <i>Cucumis melo</i>	Downy mildew	2-3 kg / ha	3 to 5 days
ONION <i>Allium cepa</i>	Downy mildew	2-3 kg / ha	3 to 5 days
LETTUCE <i>Lactuca sativa</i>	Downy mildew	2-3 kg / ha	3 to 5 days

APPLICATION Equipment

Ground equipment: Use a water volume of 400-800 L / ha with a pressure of 100-180 psi. 2 and closed cone nozzles.

Aerial equipment: Use a water volume of 80-100 L / ha.

Recommendations for the mixture:

- Make sure application equipment is thoroughly cleaned of residues of previous cargoes.
- Fill the sprayer tank with water up to a quarter of its capacity.
- Start agitation system (never with bare hands).
- Make a premix with the calculated amount of **Curzate® M-8**.
- Add the premix tank.
- We recommend the addition of water-soluble surfactants.
- Finish filling the tank with water and keep stirring.
- Use enough water to obtain full coverage of the crop.

Time re-entry into treated areas: Do not enter or allow others to enter the applied area without proper protective equipment in a span of 24 hours after application.

PHYTOTOXICITY: **Curzate® M-8** is not phytotoxic to crops of tomatoes, potatoes and tobacco in the doses recommended here, if you follow the instructions provided on the label.

INCOMPATIBILITY: **Curzate® M-8** is incompatible with alkaline products. **Curzate® M-8** is compatible with most pesticides neutral or acid reaction commonly used in tomato crops, potatoes and tobacco.

In case of poisoning, TAKE PATIENT TO DOCTOR And show the label

FIRST AID:

If Swallowed: Give 2 glasses of water immediately and induce vomiting.

If the person is unconscious, do not give anything by mouth. Call a physician.

If inhaled: Remove affected person from contaminated area and lay it in a cool and well ventilated. If the person is not breathing, give artificial respiration. If breathing is difficult, give oxygen.

If contact with eyes: Flush with water for at least 15 minutes.

If irritation persists, see a doctor.

If contact with skin: Wash affected area with soap and water for at least 15 minutes.

If irritation persists, see a doctor.

RECOMMENDATIONS TO PHYSICIAN:

Chemical group: An acetamide (Cymoxanil) plus a dithiocarbamate (mancozeb).

Symptoms: Dizziness, headache, cough, confusion, incoordination and unconsciousness. Irritation and discomfort of the skin or itching. Discomfort of the eyes, tearing and blurred vision.

Treatment: There is no specific antidote. Provide symptomatic treatment.

THE ENVIRONMENT PROTECTION MEASURES:

- This product is toxic to fish and other aquatic organisms. Do not contaminate or flowing bodies of water, food or feed by washing the application equipment, discarded or surplus store this product.

- Do not apply this product when environmental conditions favor drift or derived from the product to other areas. As the droplet size is larger, the lower the risk of drift, however, it is necessary to follow the appropriate application practices. The conditions of spray equipment, its calibration and prevent this derived from the product are the responsibility of the applicator. Consult your agricultural technician.

- Empty the contents of the container in the spray tank. Follow the local authority procedures.

CONDITIONS OF STORAGE AND TRANSPORT: Store and transport this product in a cool, dry, well ventilated and locked. Prevent product is exposed to high temperatures or below 0 ° C.

NOTICE OF WARRANTY:

The manufacturer guarantees that the contents of this package is consistent with the chemical composition indicated on the label. Not having control of the application, dosage and other factors related to its use, the manufacturer is not responsible for the results obtained as a result of misuse this product.